arXiv:1405.5619v1 [math.PR] 22 May 2014

A Characterization of Chover-Type Law of Iterated Logarithm

Deli Li* · **Pingyan Chen**† ‡

Abstract Let $0 < \alpha \leq 2$ and $-\infty < \beta < \infty$. Let $\{X_n; n \geq 1\}$ be a sequence of independent copies of a real-valued random variable X and set $S_n = X_1 + \cdots + X_n,\ n \geq 1$. We say X satisfies the (α, β)-Chover-type law of the iterated logarithm (and write $X \in CTLIL(\alpha, \beta)$) if $\limsup_{n\to\infty} \left|\frac{S_n}{n^{1/\alpha}}\right|^{(\log\log n)^{-1}} = e^{\beta}$ almost surely. This paper is devoted to a characterization of $X \in CTLIL(\alpha, \beta)$. We obtain sets of necessary and sufficient conditions for $X \in CTLIL(\alpha, \beta)$ for the five cases: $\alpha = 2$ and $0 < \beta < \infty$, $\alpha = 2$ and $\beta = 0$, $1 < \alpha < 2$ and $-\infty < \beta < \infty$, $\alpha = 1$ and $-\infty < \beta < \infty$, and $0 < \alpha < 1$ and $-\infty < \beta < \infty$. As for the case where $\alpha = 2$ and $-\infty < \beta < 0$, it is shown that $X \notin CTLIL(2, \beta)$ for any real-valued random variable X. As a special case of our results, a simple and precise characterization of the classical Chover law of the iterated logarithm (i.e., $X \in CTLIL(\alpha, 1/\alpha)$) is given; that is, $X \in CTLIL(\alpha, 1/\alpha)$ if and only if $\inf\left\{b :\ \mathbb{E}\left(\frac{|X|^{\alpha}}{(\log(e\vee|X|))^{b\alpha}}\right) < \infty\right\} = 1/\alpha$ where $\mathbb{E}X = 0$ whenever $1 < \alpha \leq 2$.

Keywords (α, β)-Chover-type law of the iterated logarithm · Sums of i.i.d. random variables · Symmetric stable distribution with exponent α

Mathematics Subject Classification (2000) Primary: 60F15; Secondary: 60G50

Running Head: Chover-type law of iterated logarithm

1 Introduction

Throughout, $\{X_n;\ n \geq 1\}$ is a sequence of independent copies of a real-valued random variable X. As usual, the partial sums of independent identically distributed (i.i.d.) random variables $X_n,\ n \geq 1$ will be denoted by $S_n = \sum_{i=1}^{n} X_i,\ n \geq 1$. Write $Lx = \log(e \vee x),\ x \geq 0$.

When X has a symmetric stable distribution with exponent $\alpha \in (0, 2)$, i.e., $\mathbb{E}\left(e^{itX}\right) = e^{-|t|^{\alpha}}$ for $t \in (-\infty, \infty)$, Chover proved that

$$\limsup_{n\to\infty} \left|\frac{S_n}{n^{1/\alpha}}\right|^{(\log\log n)^{-1}} = e^{1/\alpha} \ \text{ almost surely (a.s.)} \tag{1.1}$$

This is what we call the classical Chover law of iterated logarithm (LIL). Since then, several papers have been devoted to develop the classical Chover LIL. See, for example, Hedye [6] showed that (1.1) holds when X is in the domain of normal attraction of a nonnormal stable law, Pakshirajan and Vasudeva [8] discussed the limit points of the sequence $\{|S_n/n^{1/\alpha}|^{(\log\log n)^{-1}}; n \geq 2\}$, Kuelbs and Kurtz [7] obtained the classical Chover LIL in a Hilbert space setting, Chen [1] obtained the classical Chover LIL for the weighed sums, Vasudeva [12], Qi and Cheng [10], Peng and Qi [9] established the Chover LIL when X is in the domain of attraction of a nonnormal stable law, Scheffler [11] studied the classical Chover LIL when X is in the generalized domain of operator

*Deli Li, Department of Mathematical Sciences, Lakehead University, Thunder Bay, Ontario P7B 5E1, Canada
e-mail: dli@lakeheadu.ca

†Pingyan Chen, Department of Mathematics, Jinan University, Guangzhou 510630, China
e-mail: tchenpy@jnu.edu.cn

‡Corresponding author: Pingyan Chen (Telephone: +86 15918852098)

semistable attraction of some nonnormal law, Chen and Hu [2] extended the results of Kuelbs and Kurtz [7] to an arbitrary real separable Banach space, and so on. It should be to point out that the previous papers only gave sufficient conditions for the classical Chover LIL.

Motivated by the previous study of the classical Chover LIL, we introduce a general Chover-type LIL as follows.

Definition 1.1. *Let $0 < \alpha \leq 2$ and $-\infty < \beta < \infty$. Let $\{X, X_n;\ n \geq 1\}$ be a sequence of real-valued i.i.d. random variables. We say X satisfies the (α, β)-Chover-type law of the iterated logarithm (and write $X \in CTLIL(\alpha, \beta)$) if*

$$\limsup_{n\to\infty} \left|\frac{S_n}{n^{1/\alpha}}\right|^{(\log\log n)^{-1}} = e^{\beta} \quad a.s. \tag{1.2}$$

From the classical Chover LIL and Definition 1.1, we see that $X \in CTLIL(\alpha, 1/\alpha)$ (i.e., (1.2) holds with $\beta = 1/\alpha$) when X has a symmetric stable distribution with exponent $\alpha \in (0, 2)$.

This paper is devoted to a characterization of $X \in CTLIL(\alpha, \beta)$. The main results are stated in Section 2. We obtain sets of necessary and sufficient conditions for $X \in CTLIL(\alpha, \beta)$ for the five cases: $\alpha = 2$ and $0 < \beta < \infty$ (see Theorem 2.1), $\alpha = 2$ and $\beta = 0$ (see Theorem 2.2), $1 < \alpha < 2$ and $-\infty < \beta < \infty$ (see Theorem 2.3), $\alpha = 1$ and $-\infty < \beta < \infty$ (see Theorem 2.4), and $0 < \alpha < 1$ and $-\infty < \beta < \infty$ (see Theorem 2.5). The proofs of Theorems 2.1-2.5 are given in Section 4. For proving Theorems 2.1-2.5, three preliminary lemmas are stated in Section 3. Some llustrative examples are provided in Section 5.

2 Statement of the Main Results

The main results of this paper are the following five theorems. We begin with the case where $\alpha = 2$ and $0 < \beta < \infty$.

Theorem 2.1. *Let $0 < \beta < \infty$. Let $\{X, X_n; n \geq 1\}$ be a sequence of i.i.d. real-valued random variables. Then*

$$X \in CTLIL(2, \beta),\ \textit{i.e.,}\ \limsup_{n\to\infty} \left|\frac{S_n}{\sqrt{n}}\right|^{(\log\log n)^{-1}} = e^{\beta} \quad a.s. \tag{2.1}$$

if and only if

$$\mathbb{E}X = 0 \quad \textit{and} \quad \inf\left\{b > 0 :\ \mathbb{E}\left(\frac{X^2}{(L|X|)^{2b}}\right) < \infty\right\} = \beta. \tag{2.2}$$

For the case where $\alpha = 2$ and $\beta = 0$, we have the following result.

Theorem 2.2. *Let $\{X, X_n; n \geq 1\}$ be a sequence of i.i.d. non-degenerate real-valued random variables. Then*

$$\limsup_{n\to\infty} \left|\frac{S_n}{\sqrt{n}}\right|^{(\log\log n)^{-1}} \leq 1 \quad a.s. \tag{2.3}$$

if and only if

$$\mathbb{E}X = 0 \quad \textit{and} \quad \inf\left\{b > 0 :\ \mathbb{E}\left(\frac{X^2}{(L|X|)^{2b}}\right) < \infty\right\} = 0. \tag{2.4}$$

In either case, we have

$$X \in CTLIL(2,0),\ \textit{i.e.,}\ \limsup_{n\to\infty}\left|\frac{S_n}{\sqrt{n}}\right|^{(\log\log n)^{-1}} = 1 \quad a.s. \tag{2.5}$$

Remark 2.1. *Let c be a constant. Note that*

$$\limsup_{n\to\infty}\left|\frac{nc}{\sqrt{n}}\right|^{(\log\log n)^{-1}} = \begin{cases} 0 & if\ c = 0, \\ \infty & if\ c \neq 0. \end{cases}$$

Thus, from Theorem 2.2, we conclude that, for any $-\infty < \beta < 0$, $X \notin CTLIL(2,\beta)$ for any real-valued random variable X.

In the next three theorems, we provide necessary and sufficient conditions for $X \in CTLIL(\alpha,\beta)$ for the three cases where $1 < \alpha < 2$ and $-\infty < \beta < \infty$, $\alpha = 1$ and $-\infty < \beta < \infty$, and $0 < \alpha < 1$ and $-\infty < \beta < \infty$ respectively.

Theorem 2.3. *Let $1 < \alpha < 2$ and $-\infty < \beta < \infty$. Let $\{X, X_n; n \geq 1\}$ be a sequence of i.i.d. real-valued random variables. Then*

$$X \in CTLIL(\alpha,\beta),\ \textit{i.e.,}\ \limsup_{n\to\infty}\left|\frac{S_n}{n^{1/\alpha}}\right|^{(\log\log n)^{-1}} = e^{\beta} \quad a.s.$$

if and only if

$$\mathbb{E}X = 0 \quad \textit{and} \quad \inf\left\{b:\ \mathbb{E}\left(\frac{|X|^{\alpha}}{(L|X|)^{b\alpha}}\right) < \infty\right\} = \beta.$$

Theorem 2.4. *Let $-\infty < \beta < \infty$. Let $\{X, X_n; n \geq 1\}$ be a sequence of i.i.d. real-valued random variables. Then*

$$X \in CTLIL(1,\beta),\ \textit{i.e.,}\ \limsup_{n\to\infty}\left|\frac{S_n}{n}\right|^{(\log\log n)^{-1}} = e^{\beta} \quad a.s.$$

if and only if

$$\begin{cases} \inf\left\{b:\ \mathbb{E}\left(\dfrac{|X|}{(L|X|)^{b}}\right) < \infty\right\} = \beta & \textit{if } \beta > 0, \\ \textit{either}\ \ \mathbb{E}|X| < \infty \textit{ and } \mathbb{E}X \neq 0 \quad \textit{or} \quad \inf\left\{b:\ \mathbb{E}\left(\dfrac{|X|}{(L|X|)^{b}}\right) < \infty\right\} = 0 & \textit{if } \beta = 0, \\ \mathbb{E}X = 0 \textit{ and } \inf\left\{b:\ \mathbb{E}\left(\dfrac{|X|}{(L|X|)^{b}}\right) < \infty\right\} = \beta & \textit{if } \beta < 0. \end{cases}$$

In particular, $\mathbb{E}|X| < \infty$ and $\mathbb{E}X \neq 0$ imply that

$$\lim_{n\to\infty}\left|\frac{S_n}{n}\right|^{(\log\log n)^{-1}} = 1 \quad a.s.$$

Theorem 2.5. *Let $0 < \alpha < 1$ and $-\infty < \beta < \infty$. Let $\{X, X_n; n \geq 1\}$ be a sequence of i.i.d. real-valued random variables. Then*

$$X \in CTLIL(\alpha, \beta), \ \textit{i.e.,} \ \limsup_{n\to\infty} \left| \frac{S_n}{n^{1/\alpha}} \right|^{(\log\log n)^{-1}} = e^{\beta} \quad a.s.$$

if and only if

$$\inf\left\{ b : \ \mathbb{E}\left(\frac{|X|^{\alpha}}{(L|X|)^{b\alpha}} \right) < \infty \right\} = \beta.$$

Remark 2.2. *From our Theorems 2.1, 2.3, 2.4, and 2.5, a simple and precise characterization of the classical Chover LIL (i.e., $X \in CTLIL(\alpha, 1/\alpha)$) is obtained as follows. For $0 < \alpha \leq 2$, we have*

$$X \in CTLIL(\alpha, 1/\alpha), \ \textit{i.e.,} \ \limsup_{n\to\infty} \left| \frac{S_n}{n^{1/\alpha}} \right|^{(\log\log n)^{-1}} = e^{1/\alpha} \quad a.s.$$

if and only if

$$\inf\left\{ b : \ \mathbb{E}\left(\frac{|X|^{\alpha}}{(L|X|)^{b\alpha}} \right) < \infty \right\} = 1/\alpha \quad \textit{where} \ \ \mathbb{E}X = 0 \quad \textit{whenever} \ \ 1 < \alpha \leq 2.$$

Our Theorems 2.1-2.5 also imply the following two interesting results.

Corollary 2.1. *Let $0 < \alpha \leq 2$. Let $\{X, X_n; n \geq 1\}$ be a sequence of i.i.d. real-valued random variables. Then*

$$\lim_{n\to\infty} \left| \frac{S_n}{n^{1/\alpha}} \right|^{(\log\log n)^{-1}} = 0 \quad a.s.$$

if and only if

$$\begin{cases} \inf\left\{ b : \ \mathbb{E}\left(\dfrac{|X|^{\alpha}}{(L|X|)^{b\alpha}} \right) < \infty \right\} = -\infty & \textit{if } 0 < \alpha < 1, \\ \mathbb{E}X = 0 \ \textit{and} \ \inf\left\{ b : \ \mathbb{E}\left(\dfrac{|X|^{\alpha}}{(L|X|)^{b\alpha}} \right) < \infty \right\} = -\infty & \textit{if } 1 \leq \alpha < 2, \\ X = 0 \quad a.s. & \textit{if } \alpha = 2. \end{cases}$$

Corollary 2.2. *Let $0 < \alpha \leq 2$. Let $\{X, X_n; n \geq 1\}$ be a sequence of i.i.d. real-valued random variables. Then*

$$\limsup_{n\to\infty} \left| \frac{S_n}{n^{1/\alpha}} \right|^{(\log\log n)^{-1}} = \infty \quad a.s.$$

if and only if

$$\begin{cases} \inf\left\{ b : \ \mathbb{E}\left(\dfrac{|X|^{\alpha}}{(L|X|)^{b\alpha}} \right) < \infty \right\} = \infty & \textit{if } 0 < \alpha \leq 1, \\ \textit{eithor } \mathbb{E}X \neq 0 \ \textit{or} \ \inf\left\{ b : \ \mathbb{E}\left(\dfrac{|X|^{\alpha}}{(L|X|)^{b\alpha}} \right) < \infty \right\} = \infty & \textit{if } 1 < \alpha \leq 2. \end{cases}$$

3 Preliminary lemmas

To prove the main results, we use the following three preliminary lemmas. The first lemma is new and may be of independent interest.

Lemma 3.1. *Let $\{a_n;\ n \geq 1\}$ be a sequence of real numbers. Let $\{c_n;\ n \geq 1\}$ be a sequence of positive real numbers such that*

$$\lim_{n\to\infty} c_n = \infty. \tag{3.1}$$

Then we have

(i) There exists a constant $-\infty < \beta < \infty$ such that

$$\limsup_{n\to\infty} |a_n|^{1/c_n} = e^{\beta} \tag{3.2}$$

if and only if

$$\limsup_{n\to\infty} \frac{|a_n|}{e^{bc_n}} = \begin{cases} 0 & \text{for all } b > \beta, \\ \infty & \text{for all } b < \beta; \end{cases} \tag{3.3}$$

(ii) There exists a constant $-\infty < \beta < \infty$ such that

$$\liminf_{n\to\infty} |a_n|^{1/c_n} = e^{\beta}$$

if and only if

$$\liminf_{n\to\infty} \frac{|a_n|}{e^{bc_n}} = \begin{cases} 0 & \text{for all } b > \beta, \\ \infty & \text{for all } b < \beta; \end{cases}$$

(iii) There exists a constant $-\infty < \beta < \infty$ such that

$$\lim_{n\to\infty} |a_n|^{1/c_n} = e^{\beta}$$

if and only if

$$\lim_{n\to\infty} \frac{|a_n|}{e^{bc_n}} = \begin{cases} 0 & \text{for all } b > \beta, \\ \infty & \text{for all } b < \beta. \end{cases}$$

Proof We prove the sufficiency part of Part (i) first. It follows from (3.3) that the set

$$\left\{n \geq 1;\ \frac{|a_n|}{e^{bc_n}} > 1\right\} \begin{cases} \text{has finitely many elements} & \text{if } b > \beta, \\ \text{has infinitely many elements} & \text{if } b < \beta. \end{cases}$$

Note that

$$\left\{n \geq 1;\ |a_n|^{1/c_n} > e^b\right\} = \left\{n \geq 1;\ \frac{|a_n|}{e^{bc_n}} > 1\right\}.$$

We thus conclude that

$$\limsup_{n\to\infty} |a_n|^{1/c_n} \begin{cases} \leq e^b & \text{for all } b > \beta, \\ \geq e^b & \text{for all } b < \beta \end{cases}$$

which ensures (3.2).

We now prove the necessity part of Part (i). For all $b \neq \beta$, let $h = (b+\beta)/2$. Then

$$\begin{cases} \beta < h < b & \text{if } b > \beta, \\ b < h < \beta & \text{if } b < \beta. \end{cases}$$

It follows from (3.2) that the set

$$\left\{ n \geq 1;\ |a_n|^{1/c_n} > e^h \right\} \begin{cases} \text{has finitely many elements} & \text{if } b > \beta, \\ \text{has infinitely many elements} & \text{if } b < \beta. \end{cases}$$

Note that

$$\left\{ n \geq 1;\ \frac{|a_n|}{e^{hc_n}} > 1 \right\} = \left\{ n \geq 1;\ |a_n|^{1/c_n} > e^h \right\}.$$

We thus have that

$$\limsup_{n\to\infty} \frac{|a_n|}{e^{hc_n}} \begin{cases} \leq 1 & \text{if } b > \beta, \\ \geq 1 & \text{if } b < \beta. \end{cases} \tag{3.4}$$

Note that (3.1) implies that

$$\lim_{n\to\infty} \frac{e^{hc_n}}{e^{bc_n}} = \lim_{n\to\infty} e^{(h-b)c_n} = \begin{cases} 0 & \text{if } b > \beta, \\ \infty & \text{if } b < \beta. \end{cases}$$

Thus it follows from (3.4) that

$$\limsup_{n\to\infty} \frac{|a_n|}{e^{bc_n}} = \left(\lim_{n\to\infty} \frac{e^{hc_n}}{e^{bc_n}} \right) \left(\limsup_{n\to\infty} \frac{|a_n|}{e^{hc_n}} \right) = \begin{cases} 0 & \text{if } b > \beta, \\ \infty & \text{if } b < \beta, \end{cases}$$

i.e., (3.3) holds.

We leave the proofs of Parts (ii) and (iii) to the reader since they are similar to the proof of Part (i). This completes the proof of Lemma 3.1. $\square$

The following result is a special case of Corollary 2 of Einmahl and Li [4].

Lemma 3.2. *Let $b > 0$. Let $\{X, X_n; n \geq 1\}$ be a sequence of i.i.d. random variables. Then*

$$\lim_{n\to\infty} \frac{S_n}{\sqrt{n}(\log n)^b} = 0 \quad a.s.$$

if and only if

$$\mathbb{E}X = 0, \quad \mathbb{E}\left(\frac{X^2}{(L|X|)^{2b}} \right) < \infty, \quad \textit{and } \lim_{x\to\infty} \frac{LLx}{(Lx)^{2b}} H(x) = 0,$$

where $H(x) = \mathbb{E}\left(X^2 I\{|X| \leq x\}\right)$, $x \geq 0$.

The following result is a generalization of Kolmogorov-Marcinkiewicz-Zygmund strong law of large numbers and follows easily from Theorems 1 and 2 of Feller [5].

Lemma 3.3. *Let $0 < \alpha < 2$ and $-\infty < b < \infty$. Let $\{X, X_n; n \geq 1\}$ be a sequence of i.i.d. random variables. Then*

$$\lim_{n\to\infty} \frac{S_n}{n^{1/\alpha}(\log n)^b} = 0 \quad a.s.$$

if and only if

$$\mathbb{E}\left(\frac{|X|^\alpha}{(L|X|)^{b\alpha}}\right) < \infty$$

where $\mathbb{E}X = 0$ whenever either $1 < \alpha < 2$ or $\alpha = 1$ and $-\infty < b \leq 0$.

4 Proofs of the Main Results

In this section, we only give the proofs of Theorems 2.1-2.2. By applying Lemmas 3.1 and 3.3, the proofs of Theorems 2.3-2.5 involves only minor modifications of the proof of Theorem 2.1 and will be omitted.

Proof of Theorem 2.1 We prove the sufficiency part first. Note that the second part of (2.2) implies that

$$\mathbb{E}\left(\frac{X^2}{(L|X|)^{2b}}\right) < \infty \text{ for all } b > \beta. \tag{4.1}$$

We thus see that for all $b > \beta$,

$$\begin{aligned} H(x) &= \mathbb{E}\left(X^2 I\{|X| \leq x\}\right) \\ &\leq \mathbb{E}\left(\frac{X^2}{(L|X|)^{2h}}(Lx)^{2h} I\{|X| \leq x\}\right) \\ &\leq (Lx)^{2h}\mathbb{E}\left(\frac{X^2}{(L|X|)^{2h}}\right), \end{aligned}$$

where $h = (b+\beta)/2$. Since $\beta < h < b$, it follows from (4.1) that

$$\frac{LLx}{(Lx)^{2b}} H(x) \leq \frac{LLx}{(Lx)^{2(b-h)}}\mathbb{E}\left(\frac{X^2}{(L|X|)^{2h}}\right) \to 0 \text{ as } x \to \infty.$$

We thus conclude from (2.2) that

$$\mathbb{E}X = 0, \ \ \mathbb{E}\left(\frac{X^2}{(L|X|)^{2b}}\right) < \infty, \ \text{ and } \lim_{x\to\infty} \frac{LLx}{(Lx)^{2b}} H(x) = 0 \text{ for all } b > \beta > 0$$

which, by applying Lemma 3.2, ensures that

$$\lim_{n\to\infty} \frac{S_n}{\sqrt{n}(\log n)^b} = 0 \text{ a.s.} \quad \text{for all } b > \beta. \tag{4.2}$$

Since $\beta > 0$, the second part of (2.2) implies that

$$\mathbb{E}\left(\frac{X^2}{(L|X|)^{2b}}\right) = \infty \text{ for all } b < \beta$$

which ensures that

$$\limsup_{n\to\infty} \frac{|S_n|}{\sqrt{n}(\log n)^b} = \infty \quad \text{a.s.} \quad \text{for all } b < \beta. \tag{4.3}$$

Let $A_n = S_n/\sqrt{n}$ and $c_n = LLn$, $n \geq 1$. It then follows from (4.2) and (4.3) that

$$\limsup_{n\to\infty} \frac{|A_n|}{e^{bc_n}} = \limsup_{n\to\infty} \frac{|S_n|}{\sqrt{n}(\log n)^b} = \begin{cases} 0 \quad \text{a.s. for all } b > \beta, \\ \\ \infty \quad \text{a.s. for all } b < \beta. \end{cases} \tag{4.4}$$

By Lemma 3.1, we see that (4.4) is equivalent to

$$\limsup_{n\to\infty} |A_n|^{1/c_n} = e^{\beta} \quad \text{a.s.},$$

i.e., (2.1) holds.

We now prove the necessity part. By Lemma 3.1, (2.1) is equivalent to (4.4) which ensures that (4.2) holds. By Lemma 3.2, we conclude from (4.2) that

$$\mathbb{E}X = 0 \quad \text{and} \quad \mathbb{E}\left(\frac{X^2}{(L|X|)^{2b}}\right) < \infty \quad \text{for all } b > \beta. \tag{4.5}$$

Since $0 < \beta < \infty$, it follows from (4.5) that

$$\beta_1 \stackrel{\Delta}{=} \inf\left\{b > 0 : \ \mathbb{E}\left(\frac{X^2}{(L|X|)^{2b}}\right) < \infty\right\} \leq \beta.$$

If $\beta_1 < \beta$ then, using the argument in the proof of the sufficiency part, we have that

$$\lim_{n\to\infty} \frac{S_n}{\sqrt{n}(\log n)^b} = 0 \quad \text{a.s.} \quad \text{for all } b > \beta_1. \tag{4.6}$$

Hence, by Lemma 3.1, (4.6) implies that

$$\limsup_{n\to\infty} \left|\frac{S_n}{\sqrt{n}}\right|^{(\log\log n)^{-1}} \leq e^{\beta_1} < e^{\beta} \quad \text{a.s.}$$

which is in contradiction to (2.1). Thus (2.2) holds. The proof of Theorem 2.1 is complete. □

Proof of Theorem 2.2 Using the same argument used in the proof of the sufficiency part of Theorem 2.1, we have from (2.4) that

$$\lim_{n\to\infty} \frac{S_n}{\sqrt{n}(\log n)^b} = 0 \quad \text{a.s.} \quad \text{for all } b > 0. \tag{4.7}$$

Since X is a non-degenerate random variable, by the classical Hartman-Wintner-Strassen LIL, we have that

$$\limsup_{n\to\infty} \frac{|S_n|}{\sqrt{2n\log\log n}} > 0 \quad \text{a.s.}$$

which implies that

$$\limsup_{n\to\infty} \frac{|S_n|}{\sqrt{n}(\log n)^b} = \infty \quad \text{a.s.} \quad \text{for all } b < 0. \tag{4.8}$$

Let $A_n = S_n/\sqrt{n}$ and $c_n = LLn$, $n \geq 1$. It then follows from (4.7) and (4.8) that

$$\limsup_{n\to\infty} \frac{|A_n|}{e^{bc_n}} = \limsup_{n\to\infty} \frac{|S_n|}{\sqrt{n}(\log n)^b} = \begin{cases} 0 \ \text{ a.s. for all } b > 0, \\ \\ \infty \ \text{ a.s. for all } b < 0. \end{cases} \tag{4.9}$$

By Lemma 3.1, we see that (4.9) is equivalent to

$$\limsup_{n\to\infty} |A_n|^{1/c_n} = e^0 = 1 \ \text{ a.s.},$$

i.e., (2.5) holds, so does (2.3).

Using the same argument used in the proof of the necessity part of Theorem 2.1, we conclude from (2.3) that

$$\mathbb{E}X = 0 \ \text{ and } \ \inf\left\{b > 0 : \ \mathbb{E}\left(\frac{X^2}{(L|X|)^{2b}}\right) < \infty\right\} \leq 0.$$

Clearly

$$\inf\left\{b > 0 : \ \mathbb{E}\left(\frac{X^2}{(L|X|)^{2b}}\right) < \infty\right\} \geq 0.$$

Thus (2.4) holds. The proof of Theorem 2.2 is therefore complete. □

5 Examples

In this section, we provide the following examples to illustrate our main results. By applying Theorems 2.3-2.5, we rededuce the classical Chover LIL in the first example.

Example 5.1. *Let $0 < \alpha \leq 2$. Let X be a symmetric real-valued stable random variable with exponent α. Clearly, $\mathbb{E}X = 0$ whenever $1 < \alpha \leq 2$.*

For $0 < \alpha < 2$, we have

$$\mathbb{P}(|X| > x) \sim \left(\frac{\sin(\pi\alpha/2)\,\Gamma(\alpha)}{\pi}\right)|x|^{-\alpha} \ \ as \ \ x \to \infty,$$

it follows that

$$\mathbb{E}\left(\frac{|X|^\alpha}{(L|X|)^{b\alpha}}\right) \begin{cases} < \infty \quad if \ b > 1/\alpha, \\ \\ = \infty \quad if \ b \leq 1/\alpha \end{cases}$$

and hence that

$$\inf\left\{b : \ \mathbb{E}\left(\frac{|X|^\alpha}{(L|X|)^{b\alpha}}\right) < \infty\right\} = 1/\alpha.$$

Thus, by Theorems 2.3-2.5, $X \in CTLIL(\alpha, 1/\alpha)$ (i.e., the classical Chover LIL follows).

However, for $\alpha = 2$, we have $\mathbb{E}X^2 = 1$. Hence, by Theorems 2.1 and 2.2, we see that $X \notin CTLIL(2, 1/2)$ but $X \in CTLIL(2, 0)$.

From our second example, we will see that $X \in CTLIL(\alpha, \beta)$ for some certain α and β even if the distribution of X is not in the domain of attraction of the stable distribution with exponent α.

Example 5.2. *Let $0 < \alpha \leq 2$. Let $d_n = \exp(2^n)$, $n \geq 1$. Given $-\infty < \lambda < \infty$. Let X be a symmetric i.i.d. real-valued random variable such that*

$$\mathbb{P}(X = -d_n) = \mathbb{P}(X = d_n) = \left(\frac{c}{2}\right) \frac{\log^\lambda d_n}{d_n^\alpha} = \left(\frac{c}{2}\right) \frac{2^{n\lambda}}{d_n^\alpha}, \quad n \geq 1$$

where

$$c = c(\alpha, \lambda) = \left(\sum_{n=1}^{\infty} \frac{\log^\lambda d_n}{d_n^\alpha} \right)^{-1} > 0.$$

Then

$$\limsup_{x \to \infty} \frac{x^\alpha}{\log^\lambda x} \mathbb{P}(|X| \geq x) = c > 0 \quad \textit{and} \quad \liminf_{x \to \infty} \frac{x^\alpha}{\log^\lambda x} \mathbb{P}(|X| \geq x) = 0.$$

Thus the distribution of X is not in the domain of attraction of the stable distribution with exponent α. Also $\mathbb{E}X = 0$ whenever either $1 < \alpha \leq 2$ or $\alpha = 1$ and $\lambda < 0$. It is easy to see that

$$\mathbb{E}\left(\frac{|X|^\alpha}{(L|X|)^{b\alpha}}\right) = \left(\frac{c}{2}\right) \sum_{n=1}^{\infty} \left(2^{\lambda - b\alpha}\right)^n \begin{cases} < \infty & \textit{if } b > \lambda/\alpha, \\ = \infty & \textit{if } b \leq \lambda/\alpha \end{cases}$$

and hence that

$$\inf\left\{ b : \ \mathbb{E}\left(\frac{|X|^\alpha}{(L|X|)^{b\alpha}}\right) < \infty \right\} = \lambda/\alpha.$$

Thus, by Theorems 2.1-2.5, we have

(1) *If $\alpha = 2$ and $0 < \lambda < \infty$, then $X \in CTLIL(2, \lambda/\alpha)$.*
(2) *If $\alpha = 2$ and $-\infty < \lambda \leq 0$, then $X \in CTLIL(2, 0)$.*
(3) *If $0 < \alpha < 2$, then $X \in CTLIL(\alpha, \lambda/\alpha)$.*

Our third example shows that X may satisfy the other Chover-type LIL studied by Chen and Hu [2] when $X \notin CTLIL(\alpha, \beta)$.

Example 5.3. *Define the density function $f(x)$ of X by*

$$f(x) = \begin{cases} 0 & \textit{if } 0 \leq |x| < e, \\ \dfrac{c}{|x|^{\alpha+1}} \exp\left(p(\log|x|)^\gamma\right) & \textit{if } |x| \geq e, \end{cases}$$

where $p \neq 0$, $0 < \gamma < 1$, $c = c(\alpha, p, \gamma)$ is a positive constant such that such that $\int_{-\infty}^{\infty} f(x)dx = 1$. On simplification one can show that for any $-\infty < b < \infty$

$$\mathbb{E}\left(\frac{|X|^\alpha}{(L|X|)^{b\alpha}}\right) = 2c \int_e^\infty \frac{\exp\left(p(\log x)^\gamma\right)}{x(\log x)^{b\alpha}} dx \begin{cases} < \infty & \textit{if } p < 0, \\ = \infty & \textit{if } p > 0. \end{cases}$$

From Theorem 2.1-2.5, we have

(1) *If $\alpha = 2$ and $p < 0$, then $X \in CTLIL(2, 0)$.*
(2) *If $\alpha = 2$ and $p > 0$, then $X \notin CTLIL(2, \beta)$ for any $0 \leq \beta < \infty$.*

(3) *If* $0 < \alpha < 2$, *then* $X \notin CTLIL(\alpha, \beta)$ *for any* $-\infty < \beta < \infty$.
However, for $0 < \alpha < 2$, *by Theorem 3.1 in Chen and Hu* [2], *we have*

$$\limsup_{n\to\infty} \left|\frac{S_n}{B(n)}\right|^{(\log\log n)^{-1}} = e^{1/\alpha} \quad a.s.,$$

where $B(x)$ *is the inverse function of* $x^\alpha/\exp\left(p(\log x)^\gamma\right)$, $x \geq e$.

Acknowledgements
The research of Deli Li was partially supported by a grant from the Natural Sciences and Engineering Research Council of Canada and the research of Pingyan Chen was partially supported by the National Natural Science Foundation of China (No. 11271161)

References

[1] Chen, P.: *Limiting behavior of weighted sums with stable distributions*. Statist. Probab. Lett. **60**, 367-375 (2002).

[2] Chen, P., Hu, T.-C.: *Limiting behavior for random elements with heavy tail*. Taiwanese J. Math. **16**, 217-236 (2012).

[3] Chover, J.: *A law of the iterated logarithm for stable summands*. Proc. Amer. Math. Soc. **17**,441-443 (1966).

[4] Einmahl, U., Li, D.: *Some results on two-sided LIL behavior*. Ann. Probab. **33**, 1601-1624 (2005).

[5] Feller, W.: *A Limit Theoerm for Random Variables with Infinite Moments*. Amer. J. Math. **68**, 257-262 (1946).

[6] Heyde, C. C.: *A note concerning behaviour of iterated logarithm type*. Proc. Amer. Math. Soc. **23**, 85-90 (1969).

[7] Kuelbs, J., Kurtz, T.: *Berry-Esseen Estimates in Hilbert Space and an Application to the Law of the Iterated Logarithm*. Ann. Probab. **2**, 387-407 (1974).

[8] Pakshirajan, R. P., Vasudeva, R.: *A law of the iterated logarithm for stable summands*. Tran. Amer. Math. Soc. **232**, 33-42 (1977).

[9] Peng, L., Qi, Y.: *Chover-type laws of the iterated logarithm for weighted sums*. Statist. Probab. Lett. **65**, 401-410 (2003).

[10] Qi, Y., Cheng, P.: *On the law of the iterated logarithm for the partial sum in the domain of attraction of stable distribution*. Chinese Ann. Math. **17**(A), 195-206 (1996)(in Chinese).

[11] Scheffler, H.-P.: *A law of the iterated logarithm for heavy-tailed random vectors*. Probab. Theory. Relat. Fields, **116**, 257-271 (2000).

[12] Vasudeva, R.: *Chover's law of the iterated logarithm and weak convergence*. Acta Math. Hung. **44**, 215-221 (1984).